



Equine Reproduction Embryo Recovery Embryo Transfer

ERS is the longest, continually in existence
Equine Embryo Transfer Company
in the United States

Female Equine Embryo Survival Studied

Researchers know that many mammals can adjust the ratio of male and female young depending on the surrounding conditions at the time of conception but how precisely this is accomplished remains a matter for debate. A recent study carried out with equine embryos by Christine Aurich, DVM, PhD, and colleagues at the University of Veterinary Medicine, Vienna, Austria, has provided important information on how the survival of female embryos could be enhanced under conditions that would otherwise tend to favor the birth of males.

According to the team, because of the process involved in sperm cell formation, there should be an equal chance that a mammalian egg will be fertilized by "male" sperm, carrying a Y chromosome, as by a "female" sperm, carrying an X chromosome. The symmetry of the system ensures that roughly the same number of males and females are born, which is helpful for the species' long-term survival. Surprisingly, though, many mammals do not produce equal numbers of male and female offspring.

The discrepancy could theoretically be explained by differential fertilization efficiencies of male and female sperm (Y chromosomes are smaller than X chromosomes so perhaps male sperm can swim faster) or by different rates of survival of male and female fetuses in the uterus. It might seem as though male embryos are better able to survive under conditions of high energy intake, but how does this work?

Jana Beckelmann, a PhD candidate in Aurich's laboratory, recently published evidence that a particular protein, insulin-like growth factor-1 (IGF1) might somehow be involved. From an examination of about 30 equine embryos, Beckelmann noticed that during early pregnancy (between eight and twelve days after fertilization) the level of messenger RNA encoding IGF1 was approximately twice as high in female embryos as in male embryos.

The difference could relate to the fact that female embryos have two X chromosomes, which might produce more of a factor required for the expression of the IGF1 gene (which is not encoded on the X chromosome) than the single X chromosome in males is able to generate. Beckelmann also confirmed that the IGF1 protein was present in the embryos, confirming that the messenger RNA is actually translated to protein.

IGF1 is known to have important functions in growth and to inhibit apoptosis, or programmed cell death. As IGF1 treatment of cattle embryos produced *in vivo* (in the live horse) improves their survival, it is likely that the factor has positive effects on the development of the early embryo in the horse. So why should female embryos contain more of the factor than males?

Early pregnancy losses are unusually high in the horse and it is believed that female embryos are especially prone to spontaneous abortion. Male embryos are known to be better able to survive under high glucose concentrations, so well-nourished mares preferentially give birth to male foals.

Beckelmann says, "We think the higher IGF1 concentrations in female embryos might represent a mechanism to ensure the survival of the embryos under conditions that would otherwise strongly favor males." If this is so, the ratio of the sexes in horses could be the result of a subtle interplay between environmental and internal factors, including insulin-like growth factor-1.

The paper, "Sex-dependent insulin like growth factor-1 expression in preattachment equine embryos," has been accepted for publication in the journal *Theriogenology* and will appear in print in the issue of January 1, 2013 (Volume 79, Issue 1, 1 January 2013, pp. 193-199). The abstract is available [online](#).

*** CLICK HERE ***

Current Articles &
Research

Reproduction
Foil Disorders
Lameness

Check back often for
new additions